5836-01-MJA

Please replace all prior claims in the application with the following:

Claim 1 (currently amended): A stabilized pharmaceutical preparation, comprising: (a) an α amino acid; (b) an optional auxiliary agent for manufacturing a pharmaceutical preparation; and (c) a 4-amino-3-substituted-butanoic acid derivative, which 4-amino-3-substituted-butanoic acid derivative has the general formula:

wherein,

Sep-10-2003 06:47pm

From-

R₁ is a hydrogen atom, a hydroxyl group, a methyl group or an ethyl group; R₂ is a monovalent group selected from:

a straight or branched alkyl group of 3 - 8 carbon atoms;

a straight or branched alkylene group of 3 - 8 carbon atoms;

a straight or branched alkyl group of 3 - 8 carbon atoms which is mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 3 - 8 carbon atoms;

a cycloalkyl group of 3 - 8 carbon atoms which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

(Amendment—page 2 of 25)



a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms, is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, and one or two of the unsubstituted methylene groups (-CH₂-) are mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

(Amendment—page 3 of 25)



PATENT PFIZER ANN ARBOR MI

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, and one or two of the unsubstituted methylene groups (-CH₂-) being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, said phenyl group being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2-, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

DI

(Amendment—page 4 of 25)

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, one of the methylene groups (-CH₂-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, and one of the methylene groups (-CH₂-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)₂- and one or two of the unsubstituted methylene groups (-CH₂-) being mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

- a phenyl or naphthyl group;
- a phenyl group substituted with a methylenedioxy group;
- halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group, a phenoxy group, a phenylmethoxy group wherein said phenyl ring is mono-substituted with a halogen atom, trifluoromethyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group, a cycloalkylmethoxy group having 5 8 carbon atoms in the cycloalkyl ring, a cycloalkenylmethoxy group having 5 8 carbon atoms in the cycloalkenyl ring, a cycloalkanedienylmethoxy group having 5 8 carbon atoms in the cycloalkenyl ring, a cycloalkanedienyl ring, a cycloalkylmethoxy group wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring having 5 8 carbon atoms is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, a cycloalkenylmethoxy group wherein one of the methylene groups (-CH₂-) in said cycloalkenylmethoxy group wherein one of the methylene groups (-CH₂-) in said cycloalkenylmethoxy group wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring having 5 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, a cycloalkanedienyl-methoxy



group wherein one of the methylene groups (-CH2-) in said cycloalkanedienyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring wherein said cycloalkyl ring is mono-substituted with a halogen atom, trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH2-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)2-, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring wherein said cycloalkenyl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH2-) in said cycloalkenyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2-, or a cycloalkanedienylmethoxy group having 5 - 8 carbon atoms in the cycloalkanedienyl ring wherein said cycloalkanedienyl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH2-) in said cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2-;

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS-;

an -O-, -S- or -SS-phenyl group;

a diphenylamino group;

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a

(Amendment—page 6 of 25)



Serial No. 9/674,815 5836-01-MJA

hydroxyl group, a alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS- and mono-, dior tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an -O-, -S- or -SS-phenyl group wherein said phenyl group is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

or


R₁ and R₂, together with the carbon atom to which they are attached, may form a divalent group selected from:

a cycloalkylidene group of 5 - 8 carbon atoms;

a cycloalkylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group or a carboxyl group;

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH2-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, $-SO- or -S(O)_2-;$

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH2-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)2- group and one or more of the unsubstituted methylene groups (-CH2-) in said cycloalkyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;





a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂- group and one or more of the unsubstituted methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms, said phenyl ring being mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;



a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms; or

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

provided that when R2 is a phonyl or naphthyl group which is mone, dier tri-substituted with a halogen atom, the α amino acid is not glycine; and provided that when the 4-amino 3 substituted butanoic acid-derivative is gabapentin, the α-amino acid is not methyl D aspartic acid.

Claim 2 (previously amended): The stabilized pharmaceutical preparation of Claim 1 wherein said α-amino acid is one or more selected from

the L-, D- and DL-forms of neutral α-amino acids;

alkali salts, acid amides, alkyl-substituted derivatives of acid amides or alkyl esters of the L-, D- and DL-forms of acidic α-amino acids;

acid addition salts or monoacylated derivatives of the L-, D- and DL-forms of basic α-amino acids;

α,ω-diaminodicarboxylic acids; and

acidic amino acid-basic amino acid adducts of the L-, D- and DL-forms of acidic α -amino acids and the L-, D- and DL-forms of basic α -amino acids.



Serial No. 9/674,815 5836-01-MJA

Claim 3 (previously amended): The stabilized pharmaceutical preparation of Claim 2 wherein said α-amino acid is one or more selected from

neutral α-amino acids consisting of glycine, phenylglycine, hydroxyphenylglycine, dihydroxyphenylglycine, L-alanine, hydroxy-L-alanine, L-leucine, hydroxy-L-leucine, dihydroxy-L-leucine, L-norleucine, methylene-L-norleucine, L-ketonorleucine, Lisoleucine, hydroxy-L-isoleucine, dihydroxy-L-isoleucine, L-valine, hydroxy-L-valine, L-isovaline, L-norvaline, hydroxy-L-norvaline, hydroxy-L-ketonorvaline, L-methionine, L-homomethionine, L-ethionine, L-threonine, acetyl-L-threonine, L-tryptophan, hydroxy-L-tryptophan, methyl-L-tryptophan, L-tyrosine, hydroxy-L-tyrosine, methyl-L-tyrosine, bromo-L-tyrosine, dibromo-L-tyrosine, 3,5-diiodo-L-tyrosine, acetyl-L-tyrosine, chloro-L-tyrosine, L-m-tyrosine, L-levodopa, L-methyldopa, L-thyroxine, L-serine, acetyl-Lserine, L-homoserine, acetyl-L-homoserine, ethyl-L-homoserine, propyl-L-homoserine, butyl-L-homoserine, L-cystine, L-homocystine, methyl-L-cysteine, allyl-L-cysteine, propyl-L-cysteine, L-phenylalanine, dihydro-L-phenylalanine, hydroxymethyl-Lphenylalanine, L-aminobutyric acid, L-aminoisobutyric acid, L-ketoaminobutyric acid, dichloro-L-aminobutyric acid, dihydroxy-L-aminobutyric acid, phenyl-L-aminobutyric acid, L-aminovaleric acid, L-aminohydroxyvaleric acid, dihydroxy-L-aminovaleric acid, L-aminoisovaleric acid, L-aminohexanoic acid, methyl-L-aminohexanoic acid, Laminoheptanoic acid, L-aminooctanoic acid and citrulline and the D- and DL-forms thereof:

acidic α-amino acids consisting of L-aspartic acid, L-glutamic acid, L-carbocysteine, L-aminoglutaric acid, L-aminosuccinic acid, L-aminoadipic acid, L-aminopimelic acid, hydroxy-L-aminopimelic acid, methyl-L-aspartic acid, hydroxy-L-aspartic acid, methyl-L-glutamic acid, methyl-hydroxy-L-glutamic acid, L-methyleneglutamic acid, hydroxy-L-glutamic acid, dihydroxy-L-glutamic acid and hydroxy-L-aminoadipic acid and the D- and DL-forms thereof;

basic α-amino acids consisting of L-arginine, L-lysine, L-ornithine, L-canavanine, L-canaline, hydroxy-L-lysine, L-homoarginine, hydroxy-L-homoarginine, hydroxy-L-omithine, L-diaminopropionic acid, L-diaminopropionic aci

(Amendment—page 10 of 25)



Sep-10-2003 06:50pm

Serial No. 9/674,815 5836-01-MJA

diaminovaleric acid, L-diaminoheptanoic acid, and L-diaminooctanoic acid and the Dand DL-forms thereof; and

α,ω-diaminodicarboxylic acids consisting of diaminosuccinic acid, diaminoglutaric acid, diaminoadipic acid and diaminopimelic acid;

provided that, when said α -amino acid is an adipic α -amino acid, it is used in the form of the corresponding alkali salt, acid amide, alkyl-substituted derivative of acid amide or alkyl ester thereof, or

when said α -amino acid is a basic α -amino acid, it is used in the form of the corresponding acid addition salt or monoacylated derivative thereof, or

said acidic α -amino acid and said basic α -amino acid are also used in the form of the corresponding acidic amino acid-basic amino acid adduct.

Claim 4 (previously amended): A stabilized pharmaceutical preparation as claimed in Claims 1,

2, or 3 wherein a total amount of said α-amino acid is in the range of 0.001 - 80 moles per
mole of the 4-amino-3-substituted-butanoic acid derivative.

Claim 5 (previously amended): A stabilized pharmaceutical preparation as claimed in Claims 1, 2, 3, or 4 wherein it is in the form of liquid preparations.

Claim 6 (previously amended): A stabilized pharmaceutical preparation as claimed in Claim 5 wherein it is in the dosage form of liquid preparations, syrups or injections.

Claim 7 (previously amended): A stabilized pharmaceutical preparation as claimed in Claims 1, 2, 3, or 4 wherein it is in the form of solid preparations.

Claim 8 (previously amended): A stabilized pharmaceutical preparation as claimed in Claim 7 which is in the dosage form of tablets, powders, granules or capsules.



94

P.013/028

F-276

T-986

Claim 9 (currently amended): A stabilized pharmaceutical preparation as claimed in Claims 1, 2, 3, 4, 5, 6, 7, or 8 which is a gabapentin-containing preparation, a pregabalin-containing preparation, a baclofen-containing preparation, or a preparation containing 3-aminomethyl-4-cyclohexyl-butanoic acid, 3-aminomethyl-5-cyclohexyl-pentanoic acid, 3-aminomethyl-4-phenyl-butanoic acid or 3-aminomethyl-5-phenyl-pentanoic acid.

Claims 10-17 (cancelled)

- Claim 18 (currently amended): A stabilized pharmaceutical preparation containing an α amino acid, an auxiliary agent for manufacturing a pharmaceutical preparation if necessary, and gabapentin.
- Claim 19 (previously added): The stabilized pharmaceutical preparation of Claim 18 wherein the α amino acid is as provided in Claims 2 or 3.
- Claim 20 (currently amended): The stabilized pharmaceutical preparation containing of Claims
 18 or 19 in the form of liquid preparations, syrups, or injections.
- Claim 21 (previously added): The stabilized pharmaceutical preparation of Claims 18 or 19 in the form of solid preparations.
- Claim 22 (previously added): The stabilized pharmaceutical preparation of Claim 21 in the dosage forms of tablets, powders, granules, or capsules.
- Claim 23 (previously added): The stabilized pharmaceutical preparation of claim 1 wherein the α-amino acid is phenylglycine, hydroxyphenylglycine, dihydroxyphenylglycine, an L-α-amino acid or a DL-α-amino acid.



Serial No. 9/674,815 5836-01-MJA

Claim 24 (currently amended): The stabilized pharmacoutical preparation of claim 1 wherein the examino acid is glycine and A stabilized pharmaceutical preparation, comprising: (a) an α amino acid; (b) an auxiliary agent for manufacturing a pharmaceutical preparation; and (c) a 4-amino-3-substituted-butanoic acid derivative, which 4-amino-3-substituted-butanoic acid derivative has the general formula:

$$R_1$$
 R_2 R_2

wherein,

 R_1 is a hydrogen atom, a hydroxyl group, a methyl group or an ethyl group; R_2 is a monovalent group selected from:

a straight or branched alkyl group of 3 - 8 carbon atoms;

a straight or branched alkylene group of 3 - 8 carbon atoms;

a straight or branched alkyl group of 3 - 8 carbon atoms which is mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 3 - 8 carbon atoms;

a cycloalkyl group of 3 - 8 carbon atoms which is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 4 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an



alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms wherein said phenyl ring is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkylcycloalkyl group wherein said cycloalkyl has 3 - 8 carbon atoms, is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, and one or two of the unsubstituted methylene groups (-CH₂-) are mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;



a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, and one or two of the unsubstituted methylene groups (-CH₂-) being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkyl group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-, said phenyl group being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenyl group of 5 - 8 carbon atoms or a cycloalkanedienyl group of 5 - 8 carbon atoms, one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring being replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2-, said phenyl ring being mono- or di-substituted with a halogen atom, a

(Amendment—page 15 of 25)



trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, one of the methylene groups (-CH₂-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

an alkylcycloalkyl group wherein said cycloalkyl has 5 - 8 carbon atoms and is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-, and one of the methylene groups (-CH₂-) in said cycloalkyl ring being replaced by -O-, -NH-, -S-, -SO- or -S(O)₂- and one or two of the unsubstituted methylene groups (-CH₂-) being mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a phenyl or naphthyl group;

a phenyl group substituted with a methylenedioxy group;

a phenyl or naphthyl group which is mono-, di- or tri-substituted with a halogen, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group, a phenoxy group, a phenylmethoxy group, a phenylmethoxy group, a phenylmethoxy group, a phenylmethoxy group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring, a cycloalkanedienylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring, a cycloalkanedienyl ring, a cycloalkylmethoxy group wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring having 5 - 8 carbon atoms is replaced by



Serial No. 9/674,815 5836-01-MJA

-O-, -NH-, -S-, -SO- or -S(O)2-, a cycloalkenylmethoxy group wherein one of the methylene groups (-CH2-) in said cycloalkenyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, a cycloalkanedienyl-methoxy group wherein one of the methylene groups (-CH2-) in said cycloalkanedienyl ring having 5 - 8 carbon atoms is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2group, a cycloalkylmethoxy group having 5 - 8 carbon atoms in the cycloalkyl ring wherein said cycloalkyl ring is mono-substituted with a halogen atom, trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH2-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)2-, a cycloalkenylmethoxy group having 5 - 8 carbon atoms in the cycloalkenyl ring wherein said cycloalkenyl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxy group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH2-) in said cycloalkenyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-, or a cycloalkanedienylmethoxy group having 5 - 8 carbon atoms in the cycloalkanedienyl ring wherein said cycloalkanedienyl ring is mono-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group and one of the methylene groups (-CH2-) in said cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)2-;

an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS-;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS-;

an -O-, -S- or -SS-phenyl group;

a diphenylamino group:

(Amendment—page 17 of 25)



an alkylphenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms optionally interrupted with -O-, -S- or -SS- and mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, a alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an alkyl-O-, -S- or -SS-phenyl group wherein said phenyl group is linked to an alkylene group having 1 - 4 carbon atoms via -O-, -S- or -SS- and mono-, dior tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

an -O-, -S- or -SS-phenyl group wherein said phenyl group is mono-, di- or tri-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an amino group, a nitro group or a carboxyl group;

٥r

R₁ and R₂, together with the carbon atom to which they are attached, may form a divalent group selected from:

a cycloalkylidene group of 5 - 8 carbon atoms;

a cycloalkylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group or a carboxyl group;

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂-;

a cycloalkylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkyl ring is replaced by -O-, -NH-, -S-, -SO- or -S(O)₂- group and one or more of the unsubstituted methylene groups (-CH₂-) in said cycloalkyl ring are mono-, di-, tri- or tetra-substituted with a

(Amendment—page 18 of 25)



halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms which is mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, a cycloalkyl group, a phenyl group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂-;

a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms wherein one of the methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring is replaced by -O-, -NH-, =N-, -S-, -SO- or -S(O)₂- group and one or more of the unsubstituted methylene groups (-CH₂-) in said cycloalkenyl ring or cycloalkanedienyl ring are mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, an oxo group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkylidene group of 4 - 8 carbon atoms, said phenyl ring being mono-, di-, tri- or tetra-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl

D1

group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group;

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms; or

a condensed ring group formed by ortho-fusion of a phenyl ring with a cycloalkenylidene group of 5 - 8 carbon atoms or a cycloalkanedienylidene group of 5 - 8 carbon atoms, said phenyl ring being mono- or di-substituted with a halogen atom, a trifluoromethyl group, a hydroxyl group, an alkyl group, an alkoxy group, an alkylthio group, an amino group, a nitro group, a carboxyl group or a carboalkoxy group; and

wherein said α-amino acid is one or more selected from

neutral a-amino acids consisting of phenylglycine, hydroxyphenylglycine, dihydroxyphenylglycine, L-alanine, hydroxy-L-alanine, L-leucine, hydroxy-L-leucine, dihydroxy-L-leucine, L-norleucine, methylene-L-norleucine, L-ketonorleucine, Lisoleucine, hydroxy-L-isoleucine, dihydroxy-L-isoleucine, L-yaline, hydroxy-L-valine, <u>L-isovaline, L-norvaline, hydroxy-L-norvaline, hydroxy-L-ketonorvaline, L-methionine, </u> L-homomethionine, L-ethionine, L-threonine, acetyl-L-threonine, L-tryptophan, hydroxy-L-tryptophan, methyl-L-tryptophan, L-tyrosine, hydroxy-L-tyrosine, methyl-L-tyrosine, bromo-L-tyrosine, dibromo-L-tyrosine, 3,5-diiodo-L-tyrosine, acetyl-L-tyrosine, chloro-L-tyrosine, L-m-tyrosine, L-levodopa, L-methyldopa, L-thyroxine, L-serine, acetyl-Lserine, L-homoserine, acetyl-L-homoserine, ethyl-L-homoserine, propyl-L-homoserine, butyl-L-homoserine, L-cystine, L-homocystine, methyl-L-cysteine, allyl-L-cysteine, propyl-L-cysteine, L-phenylalanine, dihydro-L-phenylalanine, hydroxymethyl-Lphenylalanine, L-aminobutyric acid, L-aminoisobutyric acid, L-ketoaminobutyric acid, dichloro-L-aminobutyric acid, dihydroxy-L-aminobutyric acid, phenyl-L-aminobutyric acid, L-aminovaleric acid, L-aminohydroxyvaleric acid, dihydroxy-L-aminovaleric acid, L-aminoisovaleric acid, L-aminohexanoic acid, methyl-L-aminohexanoic acid, L-



aminoheptanoic acid, L-aminooctanoic acid and citrulline and the D- and DL-forms thereof;

acidic α-amino acids consisting of L-aspartic acid, L-glutamic acid, L-carbocysteine, L-aminoglutaric acid, L-aminosuccinic acid, L-aminoadipic acid, L-aminopimelic acid, hydroxy-L-aminopimelic acid, methyl-L-aspartic acid, hydroxy-L-aspartic acid, methyl-L-glutamic acid, methyl-hydroxy-L-glutamic acid, L-methyleneglutamic acid, hydroxy-L-glutamic acid, dihydroxy-L-glutamic acid and hydroxy-L-aminoadipic acid and the D- and DL-forms thereof;

basic α-amino acids consisting of L-arginine, L-lysine, L-ornithine, L-canavanine, L-canaline, hydroxy-L-lysine, L-homoarginine, hydroxy-L-homoarginine, hydroxy-L-ornithine, L-diaminopropionic acid, L-diaminohexanoic acid, L-diaminobutyric acid, L-diaminovaleric acid, L-diaminoheptanoic acid, and L-diaminooctanoic acid and the D-and DL-forms thereof; and

α,ω-diaminodicarboxylic acids consisting of diaminosuccinic acid, diaminoglutaric acid, diaminoadipic acid and diaminopimelic acid;

provided that, when said α-amino acid is an adipic α-amino acid, it is used in the form of the corresponding alkali salt, acid amide, alkyl-substituted derivative of acid amide or alkyl ester thereof, or

when said α-amino acid is a basic α-amino acid, it is used in the form of the corresponding acid addition salt or monoacylated derivative thereof, or

said acidic α-amino acid and said basic α-amino acid are also used in the form of the corresponding acidic amino acid-basic amino acid adduct.

